

C

n

Maintenance of Scienceware® Eye Wash Bottles

Catalog No. F24850-0000, F24851-0000

1. Wash the Eye Wash Bottle and all accessory parts: cap, tubes, etc. in mild detergent solution.

R

2. Rinse well to remove all traces of detergent.

S

Ν

3. Under clean conditions, fill Eye Wash Bottle completely.

(We recommend Scienceware[®] sterile eye wash solution refill # F24879-0032. Please note that once the refill bottle has been opened, the sterility of the solution cannot be assured).

- 4. Replace top, close cap and return to emergency station.
- **5.** If an eye wash bottle is used, empty it completely and proceed through steps 1-4 before replacing in station.
- **6.** After using an eye wash bottle, the injured individual should see a physician at once for examination and treatment.

To ensure cleanliness, it is advisable to establish a regular eye wash bottle cleaning and eye wash solution replacement procedure. See the refilling guidelines below.



N

S

Guidelines for Refilling Scienceware® Eye Wash Bottles

The below recommendations for solution replacement are based on information gathered from sources outside Bel-Art Products to help assist our customers. Bel-Art Products does not guarantee the accuracy of the information gathered.

Rinse solution and frequency of change:

- Water only change once a week.
- Water and Preservative (varies by manufacturer) change every two months.
- Pre-mixed Eye Wash Solution change every 4 to 6 months.
- Unopened, Sterile Eye Wash Solution (F24879-0032) until expiration date on bottle.

Bel-Art Products Sterile Eye Wash solution (F24879-0032) has a shelf life of up to 3 years - refer to expiration date on bottle. However, once the bottle has been opened, the product is no longer sterile. The Sterile Eye Wash Solution is meant for use with Scienceware[®] Eye Wash Stations (F24878-1032 and F24878-2032).

If using the Sterile Eye Wash Solution (F24879-0032) to refill Scienceware[®] Emergency Eye Wash Bottles (F24850-0000 and F24851-0000), the solution can no longer be considered sterile once transferred into those bottles. Replace transferred solution every 4 to 6 months.



PEQUANNOCK, NJ 07440-1992 USA TEL: 1-800-4BEL-ART • FAX: 973 694-7199 • www.belart.com